

NOETHER SYMMETRY APPROACH IN PURE GRAVITY WITH VARIABLE G AND Λ

Alfio Bonanno,^{1,2} Giampiero Esposito^{3,4} Claudio Rubano^{4,3} and Paolo Scudellaro^{4,3}

¹*Osservatorio Astrofisico, Via S. Sofia 78, 95123 Catania, Italy*

²*Istituto Nazionale di Fisica Nucleare, Sezione di Catania,
Corso Italia 57, 95129 Catania, Italy*

³*Istituto Nazionale di Fisica Nucleare, Sezione di Napoli,
Complesso Universitario di Monte S. Angelo,
Via Cintia, Edificio N', 80126 Napoli, Italy*

⁴*Dipartimento di Scienze Fisiche, Complesso Universitario di Monte S. Angelo,
Via Cintia, Edificio N', 80126 Napoli, Italy*

(Dated: February 7, 2008)

Abstract

We find exact cosmological solutions when the Newton parameter and the cosmological term are dynamically evolving in a renormalization-group improved Hamiltonian approach. In our derivation we use the Noether symmetry approach, leading to an interesting variable transformation which yields exact and general integration of the cosmological equations. The functional dependence of Λ on G is determined by the method itself, therefore generalizing previous results on symmetry principles in cosmology. We find new functional relations between Λ and G , jointly with power-law inflation for pure gravity.

I. INTRODUCTION

Among the most important questions in cosmology, the task of understanding the origin and the nature of the Big Bang still remains closely tied to unavoidable problems of physical description and interpretation. This is fully connected to the unexploited marriage between classical general relativity and quantum field theory. Many have been the proposals for solving the several issues arising from such an effort. An original approach suggests the possibility that the cosmological dynamics is generated by strong “renormalization group (hereafter RG) induced” quantum effects which would drive the (dimensionless) cosmological “constant” $\lambda(k)$ and Newton “constant” $g(k)$ from an ultraviolet attractive fixed point.

Within the framework of the effective average action [1, 2, 3] a suitable fixed point is in fact known to exist in the Einstein–Hilbert truncation of theory space [4, 5, 6] and in the higher-derivative generalization [7]. There are indications [5, 7, 8, 9, 10, 11, 12] that a non-Gaussian ultraviolet fixed point should indeed exist in the exact theory, implying its non-perturbative renormalizability. Within this RG-improved framework, gravitational phenomena at a typical distance scale $\ell \equiv k^{-1}$ can be described in terms of a scale-dependent effective action $\Gamma_k[g_{\mu\nu}]$, thought of as a Wilsonian coarse-grained free-energy functional which has been identified [4] with the effective average action for Euclidean quantum gravity; an exact functional RG equation for the k -dependence of Γ_k has also been derived. Non-perturbative solutions were then obtained in such a context, referred to as *quantum Einstein gravity*. The RG equations offer an explicit answer for the k -dependence of the running Newton term $G(k)$ and the running cosmological term $\Lambda(k)$, which is very important for an understanding of the Planck era immediately after the big bang and the structure of the black hole singularity [13, 14, 15]. The RG-improved Einstein equations for a homogeneous and isotropic universe are obtained, for example, identifying k with the inverse of cosmological time, $k \propto 1/t$ [13, 16] and by promoting G and Λ to dynamically evolving quantities in the Einstein equations.

This approach has been further investigated by Reuter and Weyer in [17], where an RG-improvement at the level of the action has been proposed. In particular, it has been shown that a new effective interaction of the type $\nabla_\nu G \nabla_\mu G$ should occur in the “bare” Lagrangian. It is therefore interesting to understand the Hamiltonian structure (if any) of the theory when a kinetic term involving the integral of the inner product of $\text{grad}G$ with

itself, divided by G^3 (see section 2), is present in the action. This question was addressed in [18], where it was shown that a consistent Hamiltonian structure can be obtained by considering G and Λ as dynamical variables instead of regarding them as external fields. On the other hand, it must always be checked that the solution is consistent with a realistic RG trajectory. Interestingly, there exists a class of power-law solutions for the RG-improved Arnowitt–Deser–Misner (hereafter ADM) Lagrangian which exactly reproduce the scaling law near the fixed point for which the product ΛG is constant [19].

The aim of this paper is to further extend the investigations of [18] by using the *Noether Symmetry Approach*, a simple and powerful tool to obtain general solutions of the cosmological equations [20, 21], and to check their consistency with the RG flow predicted by the β -functions for the Einstein–Hilbert Lagrangian. In particular, we shall thus show that it is possible to generate a class of cosmological solutions consistent with a scaling law of the type

$$\Lambda G^2 = \text{constant},$$

which is realized in the cross-over region close to that Gaussian fixed point, according to the phase structure described in [8, 17, 22]. The resulting cosmology can then be useful in describing the Early Universe during the Planck Era, before the classical evolution has taken over [8].

The structure of the paper is as follows: in section 2 we describe the Lagrangian formulation and find the Noether symmetry for the Lagrangian adopted to derive the RG-improved Einstein cosmological equations when the energy-momentum tensor is vanishing. Section 3 studies how this gives rise to exact and general solutions, section 4 describes the relation between Noether symmetries and RG evolution, while concluding remarks and a critical assessment are presented in sections 5 and 6.

II. LAGRANGIAN FORMULATION AND NOETHER SYMMETRY

In [18] the ADM formalism has been applied to models of gravity with variable G and Λ . The main foundational steps therein can be summarized as follows.

(i) On renormalization-group improving the gravitational Lagrangian in the ADM approach, one might think that G and Λ have the status of given external field, whose evolution is in

principle ruled by the RG flow equation. However, it is instead possible to generalize the standard ADM Lagrangian and regard G as a dynamical field obeying an Euler–Lagrange equation, the underlying idea being that all fields occurring in the Lagrangian L should be ruled by L in the first place. This makes it possible to fully exploit the potentialities of the action principle (when Λ and G are viewed as external fields one is instead halfway through towards such potentialities).

(ii) For this purpose, one adds to an action of the Einstein–Hilbert type (but with G variable so that it is brought within the integrand) two compensating terms such that the action reduces to the York–Gibbons–Hawking form for fixed G and Λ , and takes the same functional form as the ADM action for general relativity (despite having variable G and $\Lambda = \Lambda(G)$), i.e. ruled by the Lagrangian

$$L = \frac{1}{16\pi} \int \frac{N\sqrt{h}}{G} \left(K_{ij}K^{ij} - K^2 + {}^{(3)}R - 2\Lambda(G) \right) d^3x.$$

This should be supplemented by an interaction term of a kinetic type, i.e.

$$L_{\text{int}} = -\frac{\mu}{16\pi} \int \frac{g^{\rho\sigma} G_{;\rho} G_{;\sigma}}{G^3} \sqrt{-g} d^3x.$$

Non-vanishing values of μ ensure that the resulting Euler–Lagrange or Hamilton equations for G itself are well-defined and admit non-trivial solutions.

(iii) The functional relation between Λ and G : $\Lambda = \Lambda(G)$ is not an assumption but it follows directly from the RG equation. Indeed, in general terms, the RG equation reads [8]

$$\begin{aligned} \frac{\partial G}{\partial k} &= \beta_G(G(k), \Lambda(k), k), \\ \frac{\partial \Lambda}{\partial k} &= \beta_\Lambda(G(k), \Lambda(k), k), \end{aligned}$$

where β_G and β_Λ are the β -functions. Once the solution $G = G(k)$ and $\Lambda = \Lambda(k)$ is obtained, one can always eliminate the k dependence in order to obtain the trajectory $\Lambda = \Lambda(G)$. In other words, we are saying that we consider $\Lambda = \Lambda(G)$ along a given RG trajectory, i.e. a solution of the RG equation. Because of the tremendous technical difficulties we cannot actually solve the RG equation except near a fixed point, therefore we assume that a given trajectory exists, and we investigate what type of dynamics is consistent with a given $\Lambda = \Lambda(G)$ relation. Moreover, on a quite independent ground, i.e. at Hamiltonian level,

we can point out that, if Λ were independent of G , we would find from the ADM action the primary constraint of vanishing momentum π_Λ conjugate to Λ . The preservation of this primary constraint would lead to an unacceptable secondary constraint, i.e. a vanishing lapse function [18].

In a homogeneous and isotropic universe, this approach is shown to lead to power-law behaviours of the cosmic scale factor $a = a(t)$ for both pure gravity and a massless ϕ^4 theory, well in agreement with what occurs in fixed-point cosmology, once one adopts the constraint $\Lambda G = \text{const.}$ Here, we want to investigate again the pure-gravity case in homogeneous and isotropic cosmology (with a signature $-, +, +, +$ for the metric). For this purpose, let us start from the Lagrangian motivated by the previous considerations [18]

$$L = \frac{a^3}{16\pi G} \left(-6\frac{\dot{a}^2}{a^2} + \frac{6\mathcal{K}}{a^2} - 2\Lambda \right) + \frac{\mu}{16\pi} \frac{a^3 \dot{G}^2}{G^3}, \quad (2.1)$$

where $G = G(t)$, $\Lambda = \Lambda(G(t))$, $\mathcal{K} = -1, 0, 1$ (for open, spatially flat and closed universes, respectively), and $\mu \neq 0$ is an interaction parameter on which we do not have any observational constraint, since it is non-vanishing only for significant modifications of general relativity, which can indeed occur in the very early universe; dots indicate time derivatives. L can be recast in the form

$$L = \frac{1}{8\pi G} \left(-3a\dot{a}^2 + 3\mathcal{K}a - a^3\Lambda + \frac{1}{2}\mu a^3 \frac{\dot{G}^2}{G^2} \right), \quad (2.2)$$

and the resulting second-order Euler–Lagrange equations for a and G are [18]

$$\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{2a^2} + \frac{\mathcal{K}}{2a^2} - \frac{\Lambda}{2} - \frac{\dot{a}\dot{G}}{aG} + \frac{\mu\dot{G}^2}{4G^2} = 0, \quad (2.3)$$

$$\mu\ddot{G} - \frac{3}{2}\mu\frac{\dot{G}^2}{G} + 3\mu\frac{\dot{a}}{a}\dot{G} + \frac{G}{2} \left(-6\frac{\dot{a}^2}{a^2} + \frac{6\mathcal{K}}{a^2} - 2\Lambda + 2G\frac{d\Lambda}{dG} \right) = 0. \quad (2.4)$$

By virtue of the Hamiltonian constraint, we have also to consider the equation [18]

$$\frac{\dot{a}^2}{a^2} + \frac{\mathcal{K}}{a^2} - \frac{\Lambda}{3} - \frac{\mu}{6} \frac{\dot{G}^2}{G^2} = 0, \quad (2.5)$$

which can indeed be seen as equivalent to the following constraint on the *energy* function associated to L :

$$E_L \equiv \frac{\partial L}{\partial \dot{a}} \dot{a} + \frac{\partial L}{\partial \dot{G}} \dot{G} - L = 0. \quad (2.6)$$

Let us point out that this set of equations contains some undetermined parameters $(\mu, \mathcal{K})$ and, mostly important, the arbitrary function $\Lambda(G)$, which plays here a role similar to the

potential $V(\phi)$ in the case of a scalar field. This suggests the possibility of using the so-called “Noether symmetry approach” [20, 21] mentioned in the introduction in order to find a possible guess for this function. On relying upon Refs. [20, 21] for a complete discussion of this approach we limit ourselves to say that, in many circumstances (and here also as we shall see in a moment) this procedure not only allows a reduction of the dynamical system but leads to *general exact solutions*, which seems to us a major advantage. On the other hand, the physical motivation for this choice is just founded on the search for simplicity and “beauty” of the theory. Let us first show, therefore, that with a suitable choice of the function $\Lambda = \Lambda(G)$ a Noether symmetry exists for $L = L(a, G)$, i.e., for the Lagrangian viewed as a function of a and G , considered as coordinates of the associated configuration space [20, 21].

On considering the vector field

$$X \equiv \alpha(a, G) \frac{\partial}{\partial a} + \dot{\alpha} \frac{\partial}{\partial \dot{a}} + \beta(a, G) \frac{\partial}{\partial G} + \dot{\beta} \frac{\partial}{\partial \dot{G}}, \quad (2.7)$$

with $\alpha = \alpha(a, G)$ and $\beta = \beta(a, G)$ generic C^1 functions, and $\dot{\alpha} \equiv d\alpha/dt = (\partial\alpha/\partial a)\dot{a} + (\partial\alpha/\partial G)\dot{G}$, $\dot{\beta} \equiv d\beta/dt = (\partial\beta/\partial a)\dot{a} + (\partial\beta/\partial G)\dot{G}$, we impose the condition

$$\mathcal{L}_X L = 0, \quad (2.8)$$

$\mathcal{L}_X L$ being the Lie derivative of L along X . This in fact corresponds to a set of equations for $\alpha = \alpha(a, G)$, $\beta = \beta(a, G)$ and $\Lambda = \Lambda(G)$

$$G\alpha + 2aG \frac{\partial \alpha}{\partial a} - a\beta = 0, \quad (2.9)$$

$$3G\alpha - 3a\beta + 2aG \frac{\partial \beta}{\partial G} = 0, \quad (2.10)$$

$$6G^2 \frac{\partial \alpha}{\partial G} - \mu a^2 \frac{\partial \beta}{\partial a} = 0, \quad (2.11)$$

$$\frac{3\mathcal{K}}{a^2} \left(\alpha - \frac{a\beta}{G} \right) - 3\Lambda\alpha - a \frac{d\Lambda}{dG} \beta + \frac{a\Lambda\beta}{G} = 0. \quad (2.12)$$

Such a system is overdetermined, and we can in fact use only the first three equations to get the expressions of $\alpha = \alpha(a, G)$, $\beta = \beta(a, G)$ and, afterwards, use the fourth one as a consistency equation to constrain the function $\Lambda = \Lambda(G)$ and, possibly, the values of $\mathcal{K}$. As a matter of fact, a solution of Eqs. (2.9), (2.10) and (2.11) can be found by making the simple *ansatz* $\alpha \equiv A_1(a)A_2(G)$, which plugged into Eq. (2.9) constrains the form of β to be

$\beta \equiv B_1(a)B_2(G)$, too. On using also Eqs. (2.10) and (2.11), and getting rid of unnecessary integration constants, we find

$$\alpha(a, G) \equiv A_1(a)A_2(G) = a^{\frac{J}{3-2J}} G^{J-1}, \quad (2.13)$$

$$\beta(a, G) \equiv B_1(a)B_2(G) = \frac{3}{3-2J} a^{\frac{3(J-1)}{3-2J}} G^J, \quad (2.14)$$

where J is completely fixed in terms of the parameter μ that appears in the action, since

$$\mu = \frac{2}{3}(3-2J)^2. \quad (2.15)$$

From Eqs. (2.14) and (2.15), we thus get $J \neq 1, 3/2$, so that $\mu \neq 0, 2/3$. As a matter of fact, from now on we will use the J parameter instead of the μ parameter introduced in the Lagrangian L , since it makes it possible to cast the resulting formulae in a more convenient form.

As a consequence, the consistency equation (2.12) becomes

$$\frac{2JK}{(2J-3)a^2} = \frac{1}{3-2J} \left[2(1-J)\Lambda + G \frac{d\Lambda}{dG} \right] = \text{constant}, \quad (2.16)$$

since the left-hand side depends only on a while the right-hand side depends only on G . This is possible, therefore, if both

$$JK = 0 \quad (2.17)$$

and

$$2(1-J)\Lambda + G \frac{d\Lambda}{dG} = 0 \quad (2.18)$$

hold, which leads to the necessity of splitting the next considerations into two separate branches. As a matter of fact, always suitably neglecting unnecessary integration constants (except Λ_0), we find

$$X_0 \equiv X|_{J=0} = \frac{1}{G} \frac{\partial}{\partial a} - \frac{\dot{G}}{G^2} \frac{\partial}{\partial \dot{a}} + \frac{1}{a} \frac{\partial}{\partial G} - \frac{\dot{a}}{a^2} \frac{\partial}{\partial \dot{G}} \quad (2.19)$$

for $J = 0$ ($\Rightarrow \mu = 6$), any K , and

$$\Lambda = \Lambda_0 G^{-2},$$

while

$$\begin{aligned} X_J \equiv X|_{\forall J \neq 0} &= a^{\frac{J}{3-2J}} G^{J-1} \frac{\partial}{\partial a} + a G^{J-2} \left[\frac{J}{3-2J} G \dot{a} + (J-1) a^{\frac{3(J-1)}{3-2J}} \dot{G} \right] \frac{\partial}{\partial \dot{a}} \\ &+ \frac{3}{3-2J} a^{\frac{3(J-1)}{3-2J}} G^J \frac{\partial}{\partial G} + \frac{3}{3-2J} a^{\frac{5J-6}{3-2J}} G^{J-1} \left[\frac{3(J-1)}{3-2J} G \dot{a} + J a \dot{G} \right] \frac{\partial}{\partial \dot{G}} \end{aligned} \quad (2.20)$$

for $J \neq 0, 1, 3/2$ ($\Rightarrow$ any $\mu \neq 0, 2/3$), $\mathcal{K} = 0$, and $\Lambda = \Lambda_0 G^{2(J-1)}$. Let us stress that $X_J \rightarrow X_0$ for $J \rightarrow 0$, but the situation with $J = 0$ cannot indeed be treated as a mere subcase except when $\mathcal{K} = 0$ (X without any subscript stands for X_J for any J). As we said before, this result is a direct consequence of our approach. Other choices are possible. However, in the case of the scalar field in standard cosmology, we have found that the functional form of $V(\phi)$ is essentially the *only one* which allows exact integration in a simple way [23, 24]. It is unknown at present whether the same holds for the models considered in our paper, and this is a relevant topic for further research.

III. SOLUTIONS FROM NEW COORDINATES AND LAGRANGIAN

Let us now look for a change of coordinates, such that one of them is cyclic for the Lagrangian L , and the transformed Lagrangian produces equations that, now, can be easily dealt with. Since we have proved the existence of a Noether symmetry for L , in fact, we indeed expect that *there is* such a transformation $\{a, G\} \rightarrow \{u, v\}$, in which we can assume that u is the new cyclic coordinate, for instance, and try to deduce it, therefore, by solving the following system of equations:

$$i_X du = a^{\frac{J}{3-2J}} G^{J-1} \frac{\partial u}{\partial a} + \frac{3}{3-2J} a^{\frac{3(J-1)}{3-2J}} G^J \frac{\partial u}{\partial G} = 1, \quad (3.1)$$

$$i_X dv = a^{\frac{J}{3-2J}} G^{J-1} \frac{\partial v}{\partial a} + \frac{3}{3-2J} a^{\frac{3(J-1)}{3-2J}} G^J \frac{\partial v}{\partial G} = 0, \quad (3.2)$$

$i_X du$ and $i_X dv$ being the contractions between the vector field X and the differential forms du and dv , respectively [20, 21]. By virtue of the remarks made at the end of the previous section, we can treat together all the cases given by different values of J .

In order to solve this system, we can again use an *ansatz* on the forms of $u = u(a, G)$ and $v = v(a, G)$, also noting that, for our purpose, it is anyway enough to find just *one* solution. Thus, we can get rid of unnecessary constants, for example, and simply look for a particular solution. First of all, let us write Eq. (3.1) as

$$a \frac{\partial u}{\partial a} + \frac{3}{3-2J} G \frac{\partial u}{\partial G} = a^{\frac{3(1-J)}{3-2J}} G^{1-J}, \quad (3.3)$$

and let us define

$$u \equiv n a^{\frac{3(1-J)}{3-2J}} G^{1-J} + u_0, \quad (3.4)$$

n and u_0 being two generic constants. This soon solves the equation (3.3) above, fixing

$$n \equiv n(J) \equiv \frac{3 - 2J}{6(1 - J)}, \quad (3.5)$$

which must be non-vanishing, being $J \neq 3/2$. Using $v \equiv v_1(a) + v_2(G)$, on the other hand, also solves Eq. (3.2). When $J \neq 0$ (and $\mathcal{K} = 0$), apart from some integration constants and u_0 , we have (also setting $m \equiv m(J) \equiv 1 - J$, with $m \neq -1/2, 0$, so that $m \equiv 1/[2(3n - 1)]$)

$$u = u(a, G) = \frac{3 - 2J}{6(1 - J)} a^{\frac{3(1-J)}{3-2J}} G^{1-J} = n a^{\frac{1}{2n}} G^m, \quad (3.6)$$

$$v = v(a, G) = \log \left(a G^{\frac{2J-3}{3}} \right) = \log \left(a G^{-2nm} \right), \quad (3.7)$$

from which we get

$$u = u(a, G) = \frac{1}{2} a G, \quad (3.8)$$

$$v = v(a, G) = \log \left(a G^{-1} \right) \quad (3.9)$$

for $J = 0$ ($\Rightarrow n = 1/2$) and any $\mathcal{K}$.

In general, this involves

$$a = a(u, v) = \left[\frac{6(1 - J)}{3 - 2J} \right]^{\frac{3-2J}{6(1-J)}} \exp \left(\frac{v}{2} \right) u^{\frac{3-2J}{6(1-J)}} = n^{-n} \exp \left(\frac{v}{2} \right) u^n, \quad (3.10)$$

$$G = G(u, v) = \left[\frac{6(1 - J)}{3 - 2J} \right]^{\frac{1}{2(1-J)}} \exp \left(\frac{3}{2(2J - 3)} v \right) u^{\frac{1}{2(1-J)}} = \left[\frac{1}{n} \exp \left(-\frac{v}{2n} \right) u \right]^{1/(2m)}, \quad (3.11)$$

so that

$$a = a(u, v) = \sqrt{2u} \exp \left(\frac{v}{2} \right), \quad (3.12)$$

$$G = G(u, v) = \sqrt{2u} \exp \left(-\frac{v}{2} \right), \quad (3.13)$$

when $J = 0$ (in such expressions, one has $\mu = 2(3 - 2J)^2/3 = 6n^2/(1 - 3n)^2$).

We are now able to deduce the new expressions assumed by the Lagrangian L when we substitute such functions $a = a(u, v)$ and $G = G(u, v)$ into it. This in fact produces two different L' in the two separate cases we are describing, that is

$$L'_0 = -6 \exp(2v) \dot{u} \dot{v} + 3\mathcal{K} \exp(v) - \Lambda_0 \exp(3v) \quad (3.14)$$

for $J = 0$ and any $\mathcal{K}$, and

$$L'_J = -6 \exp \left(\frac{3(J - 2)}{2J - 3} v \right) \dot{u} \dot{v} - \Lambda_0 \exp(3v) = -6 \exp \left(\frac{m + 1}{2nm} v \right) \dot{u} \dot{v} - \Lambda_0 \exp(3v) \quad (3.15)$$

for $J \neq 0$ (and $\mathcal{K} = 0$). As expected, in both cases u is cyclic for L' . This means that there exists a constant of motion $\Sigma \equiv -\partial L'/\partial \dot{u}$ associated to L' , which makes it possible to solve the equations we can derive from this Lagrangian. For each case, in fact, we respectively have

$$\Sigma_0 = 6 \exp(2v) \dot{v} \quad (3.16)$$

and

$$\Sigma_J = 6 \exp\left(\frac{3(J-2)}{2J-3}v\right) \dot{v} = 6 \exp\left(\frac{m+1}{2nm}v\right) \dot{v}, \quad (3.17)$$

where $\Sigma_J \rightarrow \Sigma_0$ for $J \rightarrow 0$. As to the possible values of the constant of motion Σ , it must also be non-vanishing, since $\Sigma = 0$ soon yields $\dot{v} = 0$, which gives no dynamics, as can be seen from the expressions of L' in Eqs. (3.14) and (3.15). Moreover, it is important to stress that such new Lagrangians are non-singular, since the Hessian related to them turns out to be non-vanishing.

Last, to the new Lagrangians L' we can also associate the two *energy* functions [20, 21]

$$E'_0 = -6 \exp(2v) \dot{u} \dot{v} - 3\mathcal{K} \exp(v) + \Lambda_0 \exp(3v) \quad (3.18)$$

and

$$E'_J = -6 \exp\left(\frac{3(J-2)}{2J-3}v\right) \dot{u} \dot{v} + \Lambda_0 \exp(3v) = -6 \exp\left(\frac{m+1}{2nm}v\right) \dot{u} \dot{v} + \Lambda_0 \exp(3v), \quad (3.19)$$

respectively, so that again $E'_J \rightarrow E'_0$ when $J \rightarrow 0$ and $\mathcal{K} = 0$.

A. Case with $J = 0$ and generic $\mathcal{K}$

Equation (3.16) can be seen as a first-order differential equation for v in terms of time t (Σ_0 becoming a parameter in it), so that its general solution is

$$v = v(t) = \frac{1}{2} \log\left(\frac{\Sigma_0}{3}t + 2C_1\right), \quad (3.20)$$

where C_1 is an arbitrary integration constant. Equation (3.18) becomes therefore a first-order differential equation for u in terms of t (and Σ_0), and we find

$$u = u(t) = \frac{2}{15\sqrt{3}\Sigma_0^2}(\Sigma_0 t + 6C_1)^{3/2}(\Sigma_0 \Lambda_0 t - 15\mathcal{K} + 6C_1 \Lambda_0) + C_2, \quad (3.21)$$

C_2 being another arbitrary integration constant. From now on, we set $C_2 = 0$. This indeed makes us lose the generality of our solution, which is on the contrary guaranteed from the

existence of the three integration constants C_1 , C_2 , and Σ_0 , plus the condition $E_L = 0$. We set hereafter $C_2 = 0$ to simplify the resulting formulae.

Let us rescale time by defining $\tau \equiv \Sigma_0 t + 6C_1$ (which tells us that Σ_0 must be positive), so that

$$u(\tau) = \frac{2\tau^{3/2}(\Lambda_0\tau - 15\mathcal{K})}{15\sqrt{3}\Sigma_0^2}, \quad v(\tau) = \frac{1}{2} \log\left(\frac{\tau}{3}\right), \quad (3.22)$$

which can be substituted into Eqs. (3.8) and (3.9), eventually obtaining

$$a = a(\tau) = \frac{2\tau}{3\sqrt{5}\Sigma_0} \sqrt{\Lambda_0\tau - 15\mathcal{K}}, \quad (3.23)$$

$$G = G(\tau) = \frac{2\sqrt{\tau}}{\sqrt{15}\Sigma_0} \sqrt{\Lambda_0\tau - 15\mathcal{K}}, \quad (3.24)$$

and

$$\Lambda = \Lambda(G(\tau)) = \Lambda_0 G^{-2}(\tau) = \frac{15\Sigma_0^2 \Lambda_0}{4\tau(\Lambda_0\tau - 15\mathcal{K})}. \quad (3.25)$$

In such a cosmological setting, the Hubble parameter is

$$H = H(t) \equiv \frac{\dot{a}(t)}{a(t)} \equiv H(\tau) = \frac{\Sigma_0}{a} \frac{da}{d\tau} = \frac{3\Sigma_0}{2\tau} \frac{\Lambda_0\tau - 10\mathcal{K}}{\Lambda_0\tau - 15\mathcal{K}}. \quad (3.26)$$

Interestingly, if one studies the Hamilton equations for the pure-gravity Lagrangian (2.2) [18], and if one looks for fixed points of the resulting system, at which a, G and their conjugate momenta have vanishing time derivative, one finds again that Λ and G are related by $\Lambda = \Lambda_0 G^{-2}$ as in (3.25), but in a closed universe only.

When, in addition, we are in the spatially flat universe, with $\mathcal{K} = 0$, Eqs. (3.23), (3.24), (3.25) and (3.26) become, respectively,

$$a(\tau) = \frac{2}{3\Sigma_0} \sqrt{\frac{\Lambda_0}{5}} \tau^{3/2}, \quad G(\tau) = \frac{2}{\Sigma_0} \sqrt{\frac{\Lambda_0}{15}} \tau, \quad \Lambda(\tau) = \frac{15\Sigma_0^2}{4\tau^2}, \quad H(\tau) = \frac{3\Sigma_0}{2\tau}. \quad (3.27)$$

B. Case with $J \neq 0$ and $\mathcal{K} = 0$

Following the same steps as above, Eq. (3.17) gives

$$v = v(\tau) = \frac{2J-3}{J-2} \log\left[\frac{J-2}{2(2J-3)}\tau\right]^{1/3} = \frac{6nm}{m+1} \log\left[\frac{m+1}{12nm}\tau\right]^{1/3}, \quad (3.28)$$

because now $\tau \equiv \Sigma_J t + 6C_1$, and Eqs. (3.19) and (3.28) yield (with $C_2 = 0$)

$$u = u(\tau) = \frac{2(2J-3)\Lambda_0}{(3J-5)\Sigma_J^2} \left[\frac{J-2}{2(2J-3)}\tau\right]^{\frac{3J-5}{J-2}} = \frac{12nm\Lambda_0}{(6nm+m+1)\Sigma_J^2} \left[\frac{m+1}{12nm}\tau\right]^{\frac{6nm+m+1}{m+1}}. \quad (3.29)$$

Substituting them into Eqs. (3.10) and (3.11), we get

$$a = a(\tau) = a_0 \tau^{\frac{12n^2}{6n-1}}, \quad (3.30)$$

$$G = G(\tau) = G_0 \tau^{2(3n-1)}, \quad (3.31)$$

where we have defined

$$a_0 \equiv a_0(n) \equiv \left[\frac{12^{-\frac{6n+1}{6n-1}}}{12n-1} \left(\frac{6n-1}{n} \right)^{\frac{12n}{6n-1}} \frac{\Lambda_0}{\Sigma_J^2} \right]^n, \quad (3.32)$$

$$G_0 \equiv G_0(n) \equiv \left[\frac{(6n-1)^2 \Lambda_0}{12n^2(12n-1)\Sigma_J^2} \right]^{3n-1}. \quad (3.33)$$

Once more, we find that all expressions in this case are generalizations of the ones we have worked out above in the case with $J = 0$, with the already mentioned caution as to the terms with $\mathcal{K} \neq 0$. As a matter of fact, when $n = 1/2$ we get $12n^2/(6n-1) = 3/2$ and $a_0 = 2\sqrt{\Lambda_0}/(3\sqrt{5}\Sigma_0)$ for the scale factor $a(\tau)$, and $2(3n-1) = 1$, $G_0 = 2\sqrt{\Lambda_0}/(\sqrt{15}\Sigma_0)$ for the gravitational coupling $G(\tau)$. Also, note that in the expressions for a and G we have preferred to use the n parameter only.

There are other non-declared constraints on the allowed values of the n parameter. In fact, so far we have simply stressed that n should be non-vanishing. But, in our calculations, we have instead used more stringent constraints on its possible values. Already the introduction of m in fact gives $n \neq 1/3$; furthermore, now, we have also to set $n \neq 1/12, 1/6$ so as to obtain meaningful expressions above. As to the influence of this on the values of the interaction parameter μ occurring in the Lagrangian (2.2), it has also to be $\mu \neq 2/27$ (we already found that μ should be $\neq 0, 2/3$, by virtue of $J \neq 1, 3/2$).

IV. NOETHER SYMMETRIES AND RG EVOLUTION

In this paper we have not considered G and Λ in the RG equations as external fields, and we have therefore to check whether our solutions can represent some RG trajectory which can always be described as a given $\Lambda = \Lambda(G)$ law, as we said in the introduction. Indeed, by taking into account Eqs. (3.31) and (3.33), let us exhibit explicitly the behaviour of the function $\Lambda = \Lambda(\tau)$, which is peculiar of the pure-gravity regime of the universe when $J \neq 0$ and $\mathcal{K} = 0$:

$$\Lambda = \Lambda(\tau) = \Lambda_0 G^{2(J-1)} = \frac{12n^2(12n-1)\Sigma_J^2}{(6n-1)^2} \tau^{-2}, \quad (4.1)$$

since $J - 1 = -m = -1/[2(3n - 1)]$. This, of course, reduces to what has been found before for $J = 0$ (and $\mathcal{K} = 0$).

From (4.1) we have

$$\Lambda G^{2(1-J)} = \text{constant}, \quad (4.2)$$

therefore our solutions do not describe the evolution near the fixed point for which $\Lambda G = \text{constant}$, unless $J = 1/2$. On the other hand, it is not difficult to understand the scaling law (4.2) in terms of the phase diagram described in [22]. In fact, in the cross-over region between the non-Gaussian and the Gaussian fixed point, the dimensionful coupling constant scales approximately as a vacuum energy $\Lambda \sim k^4$, while the dimensionful Newton constant spends most of the RG-time in approaching the Gaussian fixed point. With the notation of figure 4 of [22], we are approaching the point T of the phase diagram. Therefore $G \sim k^{-2}$ and $\Lambda G^2 \sim \text{constant}$. Since we know from the RG-evolution that the anomalous dimension in this region is approximately vanishing, we conclude that only the $J \sim 0$ values are consistent with the RG evolution in this region. Other values of J can still represent viable cosmologies, but their physical meaning in terms of the RG flow is still obscure to us. On the other hand, the cosmology described by the $J=0$ solution can be thought of as an effective description of the Universe near the Planck era, before the transition to classical cosmology of Friedmann–Lemaître–Robertson–Walker type, when the seeds for cosmological perturbations are generated.

V. THE EXPONENT p AND PARAMETER μ AS FUNCTIONS OF J

In this section we begin by paying particular attention to the possibility of obtaining inflation from our model. As is well known, the problem of inflationary models is far from being solved. The use of a scalar field, together with a wide range of possible potentials and couplings, allows for good agreement with observation and theoretical requirements, but the nature of the scalar field remains completely unknown. As we said above, our approach, both for the choice of treating G as a variable, as well as for the Noether Symmetry Approach, seems to us more natural. As we show below, we obtain a power-law inflation with arbitrarily large exponent, which seems very encouraging. Of course, much work should still be done. In particular, the problem of graceful exit from inflation and that of the perturbations' spectrum should be faced, but this seems a rather difficult task.

While in [18] a power-law behaviour for the scale factor was guessed to solve the equations, this now has been obtained exactly and, in a sense, more generally from them. To better understand what we mean by this, let us first recall that, when $\mathcal{K} = 0$, in [18] it was supposed $a = At^\alpha$ for the scale factor; with arbitrary A , this gives $\alpha_\pm = (3 \pm \sqrt{9 + 12\xi^2\lambda_*})/6$, which is closely connected to the hypothesis of being in the neighbourhood of an ultraviolet fixed point (g_*, λ_*) , hence constraining the evolution of G and Λ to be given by

$$G = G(t) = g_*\xi^{-2}t^2, \quad \Lambda = \Lambda(t) = \lambda_*\xi^2t^{-2} \quad \Rightarrow \quad G\Lambda = g_*\lambda_* \equiv \text{constant}. \quad (5.1)$$

This anyway also allows arbitrarily large values for α_+ , ξ being undetermined, while constraining the possible values of the interaction parameter to be $\mu_\pm = 3\alpha_\pm/2 = (3 \pm \sqrt{9 + 12\xi^2\lambda_*})/4$ [18]. With $\lambda_* > 0$, we then have power-law inflation for the “+” sign and a possible solution of the horizon problem.

Now, to discuss what we have instead found here, let us first of all note that in our derivation we have preferred so far to restrict our attention to the particular case with $C_2 = 0$, hence simplifying the final expressions, in order to offer an easier way to discuss the results. In the case with $\mathcal{K} = 0$ we can use the generic J formulae, which also include that situation. The constant Σ_J can be set equal to 1, for example, hence arbitrarily fixing the time scale. Let us also set $C_1 = 0$ in order to get $a(0) = 0$, which is of course more delicate, even if using τ instead of t makes it less problematic. What is more important is C_2 , indeed, while we can also set $\Lambda_0 = 1$. This latter in fact appears as a factor in the expression of u , and suitably redefining C_2 we simply have a constant multiplying the scale factor a that can be fixed arbitrarily. All this changes much when $\mathcal{K} \neq 0$; in that case the above factor depends on J . Thus, without need to introduce the parameters n and m , we can look at the expressions where J occurs, just to take into account possible degenerations for some special values of J .

With $C_2 = 0$, the behaviour of the scale factor (given in Eq. (3.31) using n) also describes the general asymptotic trend of a at large τ . We find a power-law behaviour $a \sim \tau^p$ without strictly imposing to be near a fixed point. The exponent is

$$p \equiv \frac{(3 - 2J)^2}{3(J^2 - 3J + 2)}, \quad (5.2)$$

whose plot is shown in figure 1. We find inflation only when $J < 1$ or $J > 2$, p being always > 1 in these ranges; in particular, when J is near the values 1 and 2, the exponent




FIG. 1: The exponent $p = p(J)$.




FIG. 2: The parameter $\mu = \mu(J)$.

can assume arbitrarily large values, hence giving a very strong power-law behaviour. On the other hand, the interaction parameter μ is plotted in figure 2 as a function of J , hence showing that both p and μ assume symmetric values for $J < 1$ and $J > 2$. By virtue of the physical character of the μ parameter, we might think to limit our considerations to the case with $J > 2$, hence avoiding to take $J < 0$ into account. But, as we will see later, the behaviour of G is strongly affected by such a choice. In any case, there seem to exist two disconnected families of solutions, which can be probably seen as resulting from the $\dot{G}^2$ term in the equations, producing roots with different signs. As to the function $G = G(\tau)$, on setting $C_2 = 0$ we still have a power-law behaviour, with exponent

$$p' \equiv \frac{1}{1 - J}. \quad (5.3)$$

This leads to two alternative cases, one in which G increases and another in which G decreases. In any case, for $J < 1$ and $C_2 \neq 0$, G diverges as $\tau \rightarrow 0$. A special situation is

obtained when $p' = 2$, that is $J = 1/2$; this is an inflationary case, with $p = 16/9$, and we discuss it in the following section. In the $J = 0$ case, the exponent p is instead fixed to be $3/2$ (as can be seen in Eqs. (3.23) and (3.27), hence giving only a soft acceleration for the early universe.

VI. CONCLUDING REMARKS AND OPEN PROBLEMS

Even though the proximity of the fixed point is somehow partially guessed at the beginning (for physically justifying our Lagrangian formulation), we have, anyway, never used expressions like those in Eq. (5.1), but we have gained information on the $\Lambda(G)$ form by the existence of the assumed Noether symmetry. Here, we see that the choice we have consequently done on that concretely limits the generality of our solution. Thus, we discover that, when $J = 0$ and $\mathcal{K} = 0$, one has

$$G(\tau)\Lambda(\tau) = \frac{\sqrt{15\Lambda_0}\Sigma_0}{2}\tau^{-1}, \quad (6.1)$$

while, for $J \neq 0$ and $\mathcal{K} = 0$, we get

$$G(\tau)\Lambda(\tau) = \Lambda_0 G_0^{\frac{3n-2}{3n-1}} \tau^{2(3n-2)}. \quad (6.2)$$

In both cases, therefore, we do not find the typical constant behaviour for the product $G\Lambda$ in the neighbourhood of the fixed point. As a matter of fact, in Ref. [18] the behaviour we show in Eq. (4.1) is imposed from the beginning in the equations from the RG-improved Lagrangian for pure gravity, while we indeed solve them exactly by imposing a particular choice for the product $G\Lambda$ inspired by the existence of the Noether symmetry for the Lagrangian (see comments following Eq. (3.26)). It is interesting, anyway, that, in this respect, what we find in this paper can indeed be reduced to the results in Ref. [18] only when $J = 1/2$ (a single possible value, from which $n = 2/3$ and $\mu = 8/3$), hence implying $G\Lambda = \Lambda_0 \equiv g_\star \lambda_\star$. This also fixes the power-law behaviour of the scale factor to be $a(\tau) \sim \tau^{16/9}$, as said above. The fact that μ has now to be equal to $8/3$, while in Eq. (5.1) it can assume any value (the α parameter being, indeed, free to take any value), means that, even though the Noether symmetry might also be seen as a sort of peculiarity of the physical situation we have described in this paper, there is no easy extension of the solutions in Ref. [18] to ours, without a very sharp fine tuning.

As a further remark, let us anyway note that, for the scale factor behaviour, in Ref. [18] the case with $\mathcal{K} = \pm 1$ yields $\alpha = 1$, i.e., $a = At$, while we have here found (considering the situation with $J = 0$ and generic $\mathcal{K}$) the expression in Eq. (3.23). These scale factors are not in contradiction, indeed, even if the one we have here calculated contains explicitly the $\mathcal{K}$ contribution. In fact, by virtue of Eqs. (3.24) and (3.25) we get

$$G(\tau)\Lambda(\tau) = \frac{\sqrt{15}\Sigma_0\Lambda_0}{2\sqrt{\tau(\Lambda_0\tau - 15\mathcal{K})}}. \quad (6.3)$$

All these considerations seem to support the idea that the model discussed in this paper should indeed concern an era just subsequent to the one characterized by the ultraviolet fixed point we talked about at the beginning, an era in which matter fields (and other non-gravitational fields) can still be neglected with respect to the gravity contribution. At that time, then, the time variation of G and Λ were as relevant for the cosmic evolution as they were in the earlier stage, whose description has been given elsewhere (see Ref. [18] and references therein). One of the important open issues concerns, therefore, the way a link might be found between such two different physical situations, which demands much more future work.

On the other hand, our new results about the functional relation between Λ and G , as in (4.2), jointly with the power-law inflationary solutions that we have described so far, look very promising on the path towards further insight. Encouraging evidence is therefore emerging in favour of models with variable G and Λ being able to lead to a deeper understanding of modern cosmology [25, 26, 27, 28, 29].

Acknowledgments

The authors are grateful to the INFN for financial support. The work of G. Esposito has been partially supported by PRIN *SINTESI*.

-
- [1] Berges J, Tetradis N and Wetterich C 2002 *Phys. Rep.* **363** 223
 - [2] Wetterich C 2001 *Int. J. Mod. Phys. A* **16** 1951
 - [3] Reuter M and Wetterich C 1994 *Nucl. Phys. B* **427** 291
 - [4] Reuter M 1998 *Phys. Rev. D* **57** 971

- [5] Lauscher O and Reuter M 2002 *Phys. Rev. D* **65** 025013
- [6] Souma W 1999 *Prog. Theor. Phys.* **102** 181
- [7] Lauscher O and Reuter M 2002 *Class. Quantum Grav.* **19** 483
- [8] Reuter M and Saueressig F 2002 *Phys. Rev. D* **65** 065016
- [9] Bonanno A and Reuter M 2005 *J. High Energy Phys.* JHEP02(2005)035
- [10] Niedermaier M 2003 *Nucl. Phys. B* **673** 131
- [11] Niedermaier M 2002 *J. High Energy Phys.* JHEP12(2002)066
- [12] Forgacs P and Niedermayer M 2002 hep-th/0207028
- [13] Bonanno A and Reuter M 2002 *Phys. Rev. D* **65** 043508
- [14] Bonanno A and Reuter M 1999 *Phys. Rev. D* **60** 084011
- [15] Bonanno A and Reuter M 2000 *Phys. Rev. D* **62** 043008
- [16] Bonanno A and Reuter M 2002 *Phys. Lett. B* **527** 9
- [17] Reuter M and Weyer H 2004 *Phys. Rev. D* **69** 104022; *Phys. Rev. D* **70** 124028
- [18] Bonanno A, Esposito G and Rubano C 2004 *Class. Quantum Grav.* **21** 5005
- [19] Tsamis N C and Woodard R P 1993 *Phys. Lett. B* **301** 351
- [20] de Ritis R, Marmo G, Platania G, Rubano C, Scudellaro P, and Stornaiolo C 1990 *Phys. Rev. D* **42** 1091
- [21] Capozziello S, de Ritis R, Rubano C, and Scudellaro P 1996 *Rivista del Nuovo Cimento* **19(4)** 1
- [22] Reuter M and Weyer H 2004 *J. Cosmol. Astropart. Phys.* JCAP12(2004)001
- [23] de Ritis R, Marmo G, Platania G, Rubano C, Scudellaro P and Stornaiolo C 1990 *Phys. Lett. A* **149** 79
- [24] de Ritis R, Rubano C and Scudellaro P 1995 *Europhys. Lett.* **32** 185
- [25] Barvinsky A O, Kamenshchik A Yu and Karmazin I P 1993 *Phys. Rev. D* **48** 3677
- [26] Bauer F 2005 gr-qc/0512007
- [27] Lauscher O and Reuter M 2005 hep-th/0511260
- [28] Bonanno A, Esposito G and Rubano C 2003 *Gen. Rel. Grav.* **35** 1899
- [29] Bonanno A, Esposito G, Rubano C and Scudellaro P 2006 astro-ph/0612091